

The solution of the Minkowski problem for closed surfaces in Riemannian space.

Andrei I. Bodrenko ¹

Abstract

Author of this article created for the first time the method for finding solutions of the Minkowski problem for closed surfaces in Riemannian space.

Introduction

Author resolves the Minkowski problem as the problem of construction the continuous G-deformations preserving the product of principal curvatures for every point of surface in Riemannian space. G-deformation transfers every normal vector of surface in parallel along the path of the translation for each point of surface.

The Minkowski problem (MP) asks the fundamental question of differential geometry. H. Minkowski, in 1903, published first article on this subject. Generalizations of the MP in Euclidean and pseudo-Euclidean spaces were made basically by developing the methods created by H. Minkowski, A.D. Aleksandrov, A.V. Pogorelov, W.J. Firey. But nobody has studied this problem in Riemannian space.

The MP in Riemannian space is much more complicate than the MP in Euclidean space because well known methods that are useful for Euclidean and pseudo-Euclidean spaces are not valid for Riemannian space.

Author uses term of G -deformation because for surface in Riemannian space there is no such term as spherical image.

§1. Basic definitions. Statement of the main result.

Let R^3 be the three-dimensional Riemannian space with metric tensor $\tilde{a}_{\alpha\beta}$, F be the two-dimensional simply connected oriented closed surface in R^3 .

Let $F \in C^{m,\nu}$, $\nu \in (0; 1)$, $m \geq 4$. Let F has all strictly positive principal curvatures k_1 and k_2 . Let F be oriented so that mean curvature H is strictly positive. Denote $K = k_1 k_2$.

Let F be glued from the two-dimensional simply connected oriented surfaces F^+ and F^- of class $C^{m,\nu}$. Let F^+ be attached to F^- along the common boundary ∂F of class $C^{m+1,\nu}$.

¹©Andrei I. Bodrenko, associate professor, Department of Mathematics,
Volgograd State University, University Prospekt 100, Volgograd, 400062, RUSSIA.
E.-mail: bodrenko@mail.ru <http://www.bodrenko.com>

Let F^+ and F^- be given by immersions of the domain $D \subset E^2$ into R^3 by the equation: $y^\sigma = f^{\sigma\pm}(x), x \in D, f^\pm : D \rightarrow R^3$. Denote by $d\sigma(x) = \sqrt{g}dx^1 \wedge dx^2$ the area element of the surface F . We identify the points of immersion of surface F with the corresponding coordinate sets in R^3 . Without loss of generality we assume that D is unit disk. Let x^1, x^2 be the Cartesian coordinates.

Symbol $_{,i}$ denotes covariant derivative in metric of surface F . Symbol ∂_i denotes partial derivative by variable x^i . We will assume $\dot{f} \equiv \frac{df}{dt}$. We define $\Delta(f) \equiv f(t) - f(0)$. Let g_{ij} and b_{ij} be the coefficients of the first and the second fundamental form respectively.

We consider continuous deformation of the surface $F: \{F_t\}$ defined by the equations

$$y_t^\sigma = y^\sigma + z^\sigma(t), z^\sigma(0) \equiv 0, t \in [0; t_0], t_0 > 0. \quad (1.1)$$

Definition 1 . Deformation $\{F_t\}$ is called the continuous deformation preserving the product of principal curvatures (or M -deformation [21]) if the following condition holds: $\Delta(K) = 0$ and $z^\sigma(t)$ is continuous by t .

The deformation $\{F_t\}$ generates the following set of paths in R^3

$$u^{\alpha_0}(\tau) = (y^{\alpha_0} + z^{\alpha_0}(\tau)), \quad (1.2)$$

where $z^{\alpha_0}(0) \equiv 0, \tau \in [0; t], t \in [0; t_0], t_0 > 0$.

Definition 2 . The deformation $\{F_t\}$ is called the G -deformation if every normal vector of surface transfers in parallel along the path of the translation for each point of surface.

Theorem 1 . Let $F \in C^{m,\nu}, \nu \in (0; 1), m \geq 4$, be closed surface. Let F be glued from the two-dimensional simply connected oriented surfaces F^+ and F^- of class $C^{m,\nu}$. Let F^+ be attached to F^- along the common boundary ∂F of class $C^{m+1,\nu}$. Let $\tilde{a}_{\alpha\beta} \in C^{m,\nu}, \exists M_0 = \text{const} > 0$ such that $\|\tilde{a}_{\alpha\beta}\|_{m,\nu} < M_0, \|\partial\tilde{a}_{\alpha\beta}\|_{m,\nu} < M_0, \|\partial^2\tilde{a}_{\alpha\beta}\|_{m,\nu} < M_0$.

1) Then there exists $t_0 > 0$ such that for all $t \in [0, t_0)$ there exists three-parametric MG -deformation of class $C^{m-2,\nu}$ continuous by t .

2) If, at the point $T_0 \in F^+$, the following additional condition holds: $\forall t : z^\sigma(t) \equiv 0$. Then there exists $t_0 > 0$ such that for all $t \in [0, t_0)$ there exists only zero MG -deformation of class $C^{m-2,\nu}$ continuous by t .

§2. Deduction of G -deformation formulas for surfaces in Riemannian space.

We denote:

$$z^\sigma(t) = a^j(t)y^\sigma_{,j} + c(t)n^\sigma, \quad (2.1)$$

where $a^j(0) \equiv 0, c(0) \equiv 0, n^\sigma$ is unit normal vector of surface at the point (y^σ) . Therefore the deformation of surface F is defined by functions a^j and c .

We use designations from [30]. Notice that the equations of G -deformation obtained in §2 of [30] are :

$$a^l b_{li} + (1 + N_0(t, 0))c_{,i} + \partial_i a^j N_j(t, 0) + Q_i(t, 0) = 0, i = 1, 2. \quad (2.1)$$

The estimations of norms for functions $N_0(t, 0), N_j(t, 0), Q_i(t, 0)$ are derived in §3 of article [30] where the explicit forms of $N_0(t, 0), N_j(t, 0), Q_i(t, 0)$ are found.

We introduce conjugate isothermal coordinate system where $b_{ii} = V, i = 1, 2, b_{12} = b_{21} = 0$. Then we have the equation system from (2.1):

$$\begin{aligned} c_{,1}(1 + N_0) + Va^1 + N_k \partial_1 a^k + Q_1 &= 0 \\ c_{,2}(1 + N_0) + Va^2 + N_k \partial_2 a^k + Q_2 &= 0 \end{aligned} \quad (2.2)$$

We find solution of the equation system (2.2) by finding function $\dot{c}$ on functions $\dot{a}^i$.

We denote

$$\Psi_1 = -(c_{,1} \partial_2 N_0 - c_{,2} \partial_1 N_0 + \partial_1 a^k \partial_2 N_k - \partial_2 a^k \partial_1 N_k + \partial_2 Q_1 - \partial_1 Q_2)/V. \quad (2.3)$$

Then, from (2.2) and (2.3), we have the following equation (see [30]):

$$\partial_2 \dot{a}^1 - \partial_1 \dot{a}^2 + p_k \dot{a}^k = \dot{\Psi}_1, \quad (2.4)$$

where $p_1 = \partial_2(\ln V), p_2 = -\partial_1(\ln V)$. Note that p_k do not depend on t .

At first, we will solve the equation system (2.2) for surface F^+ for the case 2) of theorem 1, assuming that functions a^1 and a^2 are given. The solution of the equation system (2.2) was presented in [30].

From [30] we have that every pair of functions $\dot{a}^i \in C^{m-2,\nu}$ corresponds to the unique function $\dot{c} \in C^{m-2,\nu}$ and therefore to the unique function $c \in C^{m-2,\nu}$.

We find function $\dot{c}$ from the equation:

$$\dot{c} = L_a(\dot{c}) + \gamma_t, \quad (2.5)$$

by the method of successive approximations. Notice that L_a and γ_t were presented in [30] and have explicit forms.

In [30] was proved the following lemma for the case 2) of theorem 1:

Lemma 5.1. ([30]) *Let the following conditions hold:*

1) *metric tensor in R^3 satisfies the conditions: $\exists M_0 = \text{const} > 0$ such that $\|\tilde{a}_{\alpha\beta}\|_{m,\nu} < M_0, \|\partial \tilde{a}_{\alpha\beta}\|_{m,\nu} < M_0, \|\partial^2 \tilde{a}_{\alpha\beta}\|_{m,\nu} < M_0$.*

2) *$\exists t_0 > 0$ such that $a^k(t), \partial_i a^k(t), \dot{a}^k(t), \partial_i \dot{a}^k(t)$ are continuous by $t, \forall t \in [0, t_0], a^k(0) \equiv 0, \partial_i a^k(0) \equiv 0$.*

3) *$\exists t_0 > 0$ such that $a^i(t) \in C^{m-2,\nu}, \partial_k a^i(t) \in C^{m-3,\nu}, \forall t \in [0, t_0]$.*

Then $\exists t_* > 0$ such that the equation $\dot{c} = L_a(\dot{c}) + \gamma_t \forall t \in [0, t_*]$ has unique solution of class $C^{m-2,\nu}$ continuous by t .

Proof of lemma 5.1. was presented in [30].

Corollary.([30]) *Let the conditions of lemma 5.1. hold. Then the function $\dot{c}$ takes the form:*

$$\dot{c}(x^1, x^2, t) = \int_{(x_{(0)}^1, x_{(0)}^2)}^{(x^1, x^2)} \left(-V \dot{a}^1 \right) d\tilde{x}^1 + \left(-V \dot{a}^2 \right) d\tilde{x}^2 + P(\dot{a}^1, \dot{a}^2),$$

and for P the following inequality holds:

$$\|P(\dot{a}_{(1)}^1, \dot{a}_{(1)}^2) - P(\dot{a}_{(2)}^1, \dot{a}_{(2)}^2)\|_{m-2, \nu} \leq K_8(t)(\|\dot{a}_{(1)}^1 - \dot{a}_{(2)}^1\|_{m-2, \nu} + \|\dot{a}_{(1)}^2 - \dot{a}_{(2)}^2\|_{m-2, \nu}),$$

where for any $\varepsilon > 0$ there exists $t_0 > 0$ such that for all $t \in [0, t_0]$ the following inequality holds: $K_8(t) < \varepsilon$.

The proof follows from construction of function $\dot{c}$ (see [30]).

Note. For surface F^- we use similar methods of finding solutions of the equation system (2.2). But formula (5.6) from [30] takes the following form

$$\begin{aligned} \dot{c}(x^1, x^2, t) = & \int_{(x_{(0)}^1, x_{(0)}^2)}^{(x^1, x^2)} \left(-\frac{-V \dot{a}^1 N_0 + N_k \partial_1 \dot{a}^k + \dot{N}_k \partial_1 a^k + \dot{Q}_1}{(1 + N_0)} + \frac{\dot{N}_0(V a^1 + N_k \partial_1 a^k + Q_1)}{(1 + N_0)^2} \right) d\tilde{x}^1 + \\ & \left(-\frac{-V \dot{a}^2 N_0 + N_k \partial_2 \dot{a}^k + \dot{N}_k \partial_2 a^k + \dot{Q}_2}{(1 + N_0)} + \frac{\dot{N}_0(V a^2 + N_k \partial_2 a^k + Q_2)}{(1 + N_0)^2} \right) d\tilde{x}^2 + \\ & \int_{(x_{(0)}^1, x_{(0)}^2)}^{(x^1, x^2)} \left(-V \dot{a}^1 \right) d\tilde{x}^1 + \left(-V \dot{a}^2 \right) d\tilde{x}^2 + B_1^-, \end{aligned} \quad (2.6)$$

where B_1^- is arbitrary real parameter. B_1^- is found from the condition $\dot{c}^+ = \dot{c}^-$ on ∂F .

§3. The equations of MG -deformations.

Deduction the formulas of deformations preserving the product of principal curvatures was presented in [30]. The equation of deformations preserving the product of principal curvatures takes the following form:

$$\partial_1 \dot{a}^1 + \partial_2 \dot{a}^2 + q_k^{(b)} \dot{a}^k = \dot{\Psi}_2^{(b)}, \quad (3.1)$$

where $\dot{\Psi}_2^{(b)} = q_0^{(b)} \dot{c} - P_0(\dot{a}^1, \dot{a}^2, \partial_i \dot{a}^j)$. P_0 has explicit form. Notice that $q_k^{(b)} \in C^{m-3, \nu}$, $q_0^{(b)} \in C^{m-3, \nu}$ and do not depend on t .

We will use the following lemma form [30].

Lemma 6.2.1. ([30]) *Let the following conditions hold:*

1) *metric tensor in R^3 satisfies the conditions: $\exists M_0 = \text{const} > 0$ such that $\|\tilde{a}_{\alpha\beta}\|_{m, \nu} < M_0$, $\|\partial \tilde{a}_{\alpha\beta}\|_{m, \nu} < M_0$, $\|\partial^2 \tilde{a}_{\alpha\beta}\|_{m, \nu} < M_0$.*

2) $\exists t_0 > 0$ such that $a^k(t), \partial_i a^k(t), \dot{a}^k(t), \partial_i \dot{a}^k(t)$ are continuous by $t, \forall t \in [0, t_0], a^k(0) \equiv 0, \partial_i a^k(0) \equiv 0$.

3) $\exists t_0 > 0$ such that $a^i(t) \in C^{m-2, \nu}, \partial_k a^i(t) \in C^{m-3, \nu}, \forall t \in [0, t_0]$.

Then $\exists t_* > 0$ such that for all $t \in [0, t_*)$ $P_0 \in C^{m-3, \nu}$ and the following inequality holds:

$$\|P_0(\dot{a}_{(1)}^1, \dot{a}_{(1)}^2) - P_0(\dot{a}_{(2)}^1, \dot{a}_{(2)}^2)\|_{m-2, \nu} \leq K_9(t)(\|\dot{a}_{(1)}^1 - \dot{a}_{(2)}^1\|_{m-1, \nu} + \|\dot{a}_{(1)}^2 - \dot{a}_{(2)}^2\|_{m-1, \nu}),$$

where for any $\varepsilon > 0$ there exists $t_0 > 0$ such that for all $t \in [0, t_0)$ the following inequality holds: $K_9(t) < \varepsilon$.

The proof was presented in [30].

We observe all designations for functions and terms presented in [30]. We use estimations of norms for obtained functions from §2, §3, §7 of [30]. Formulas of functions $\dot{W}_1, \dot{W}_2, \dot{\Psi}_2$, the estimations for norms of these functions are presented in §8 of [30]. From article [30], we also use the following lemmas: 2.1, 2.2, 2.3, 3.1, 3.2, 5.1, 6.2.1, 7.1, 7.2, 7.3, 8.3.1, 8.3.2, and theorems: 1, 9.1, 9.2.

We have the following equation system of elliptic type from [30]:

$$\begin{aligned} \partial_2 \dot{a}^1 - \partial_1 \dot{a}^2 + p_k \dot{a}^k &= \dot{\Psi}_1, \\ \partial_1 \dot{a}^1 + \partial_2 \dot{a}^2 + q_k^{(b)} \dot{a}^k &= \dot{\Psi}_2^{(b)}, \end{aligned} \quad (3.2)$$

where we use (3.1). $\dot{\Psi}_2^{(b)} = q_0^{(b)} \dot{c} - P_0$. Note that $q_k^{(b)}$ do not depend on t .

Then we finally have the form of desired equation from [30]:

$$\partial_{\bar{z}} \dot{w} + A \dot{w} + B \bar{\dot{w}} + E(\dot{w}) = \dot{\Psi}. \quad (3.3)$$

Let, along the ∂F , be given vector field tangent to F . We denote it by the following formula:

$$v^\alpha = l^i y_{,i}^\alpha. \quad (3.4)$$

We consider the boundary-value condition:

$$\tilde{a}_{\alpha\beta} z^\alpha v^\beta = \tilde{\gamma}(s, t), s \in \partial D. \quad (3.5)$$

Define: $\tilde{\lambda}_k = \tilde{a}_{\alpha\beta} y_{,k}^\alpha v^\beta, k = 1, 2$.

Then boundary condition takes the form: $Re\{(a^1 + ia^2)(\tilde{\lambda}_1 - i\tilde{\lambda}_2)\} = \tilde{\gamma}$ on ∂F .

Denote: $\lambda_k = \frac{\tilde{\lambda}_k}{(\lambda_1)^2 + (\lambda_2)^2}, k = 1, 2. \dot{\varphi} = \frac{\dot{\tilde{\gamma}}}{(\lambda_1)^2 + (\lambda_2)^2}$.

Then boundary-value condition takes the form: $Re\{\bar{\lambda} \dot{w}\} = \dot{\varphi}$ on ∂F , where $|\lambda| = 1$.

We analyze the decidability of the following equation (A):

$$\partial_{\bar{z}} \dot{w} + A \dot{w} + B \bar{\dot{w}} + E(\dot{w}) = \dot{\Psi}, \quad Re\{\bar{\lambda} \dot{w}\} = \dot{\varphi} \quad on \quad \partial D, \quad (3.6)$$

$\lambda = \lambda_1 + i\lambda_2, |\lambda| \equiv 1, \lambda, \dot{\varphi} \in C^{m-2, \nu}(\partial D)$.

We will use the fact that $\dot{\Psi} = \dot{\Psi}(\dot{w}, z, t), E(\dot{w}) = E(\dot{w}, z, t), \dot{w} = \dot{w}(t), \dot{\varphi} = \dot{\varphi}(s, t), s \in \partial D, \lambda = \lambda(s), s \in \partial D$.

Let n be index of obtained boundary-value problem

$$n = \frac{1}{2\pi} \Delta_{\partial D} \arg \lambda(s). \quad (3.7)$$

Since v^α is not tangential to ∂F vector field then the index of boundary-value problem $n = 1$.

§4. Proof of theorem 1.

Proof of theorem 1.

At first, we will prove the case 2) of theorem 1.

Notice that we consider closed surface. We assume that surface consists of two surfaces F^+ and F^- with the same boundary ∂F . Let F^+ and F^- be glued. According to [17, 18], we introduce conjugate isothermal coordinate system on F . Therefore F is mapped gomeomorphically onto the domains $G^+ \subset E^2$ and $G^- \subset E^2$. G^+ and G^- have the same boundary Γ and G^- is infinite domain. Notice that Γ is simple closed smooth curve of class $C^{m+1,\nu}$. Note that ∂F is mapped gomeomorphically onto the curve Γ twice. According to [17, 18], without loss of generality we assume that for each point $M \in \partial F$ there exists unique point $s_M \in \Gamma$.

It is possible to introduce uniform conjugate isothermal coordinate system on F that maps F gomeomorphically onto the whole E^2 if we do not fix domains G^+ and G^- (see [17, 18]).

Along the ∂F , we consider the condition of contiguity of surfaces F^+ and F^- . Let $s^\alpha = s^k y_{,k}^\alpha$ be unit tangential to the ∂F vector field (see [28]). Let v^α be unit vector field which, on ∂F , satisfies the following conditions:

$$v^\alpha = l^i y_{,i}^\alpha, \quad \tilde{a}_{\alpha\beta} v^\alpha s^\beta = 0, \quad \tilde{a}_{\alpha\beta} v^\alpha v^\beta = 1.$$

Denote: $z_s^{+\alpha} = \tilde{a}_{\alpha\beta} z^{+\alpha} s^\beta$, $z_s^{-\alpha} = \tilde{a}_{\alpha\beta} z^{-\alpha} s^\beta$, $z_v^{+\alpha} = \tilde{a}_{\alpha\beta} z^{+\alpha} v^\beta$, $z_v^{-\alpha} = \tilde{a}_{\alpha\beta} z^{-\alpha} v^\beta$.

Therefore, according to [17,18], if the surfaces F^+ and F^- compose the smooth surface F then the condition of contiguity of surfaces F^+ and F^- takes the form:

$$z_s^{+\alpha} = z_s^{-\alpha}, \quad z_v^{+\alpha} = z_v^{-\alpha}, \quad c^+ = c^- \quad \text{on} \quad \partial F.$$

We have the following conditions:

$$\begin{aligned} \tilde{a}_{\alpha\beta} v^\alpha s^\beta &= 0, \\ \tilde{a}_{\alpha\beta} v^\alpha v^\beta &= 1, \\ \tilde{a}_{\alpha\beta} s^\alpha s^\beta &= 1. \end{aligned}$$

Therefore we get the following equation system:

$$g_{ik} l^i s^k = 0.$$

$$g_{ik}l^i l^k = 1.$$

$$g_{ik}s^i s^k = 1.$$

Consider the formula:

$$z^\alpha = a^i g_{,i}^\alpha + c n^\alpha.$$

Then

$$z_v = g_{ik} a^i l^k,$$

$$z_s = g_{ik} a^i s^k.$$

Therefore we have the following boundary-value condition on ∂F :

$$a^{+i} = a^{-i}.$$

We obtain on ∂F :

$$\dot{a}^{+i} = \dot{a}^{-i}.$$

In the article [30] author analyzed the decidability of the following boundary-value problem (A):

$$\partial_{\bar{z}} \dot{w} + A \dot{w} + B \bar{\dot{w}} + E(\dot{w}) = \dot{\Psi}, \quad \operatorname{Re}\{\bar{\lambda} \dot{w}\} = \dot{\varphi} \quad \text{on} \quad \partial D, \quad (4.1)$$

$|\lambda| \equiv 1$, $\lambda = \lambda_1 + i\lambda_2$, $\lambda, \dot{\varphi} \in C^{m-2,\nu}(\partial D)$.

Therefore we have the following boundary-value problem (A'):

$$\partial_{\bar{z}} \dot{w} + A \dot{w} + B \bar{\dot{w}} + E(\dot{w}) = \dot{\Psi}, \quad (\text{in } G^+ \text{ and } G^-),$$

$$\dot{a}^{+i} = \dot{a}^{-i}, \quad \text{on } \Gamma. \quad (4.2)$$

The boundary-value problem (A') takes the form:

$$\partial_{\bar{z}} \dot{w} + A \dot{w} + B \bar{\dot{w}} + E(\dot{w}) = \dot{\Psi}, \quad (\text{in } G^+ \text{ and } G^-),$$

$$\dot{w}^+ = \dot{w}^-, \quad \text{on } \Gamma. \quad (4.3)$$

The above problem is generalized Carleman problem. Since $F \in C^{m,\nu}$, $m \geq 4$ then the condition of contiguity of surfaces F^+ and F^- is of at least 4-th order.

Consider the following problem

$$\partial_{\bar{z}} \dot{w} + A \dot{w} + B \bar{\dot{w}} = 0, \quad (\text{in } G^+ \text{ and } G^-),$$

$$\dot{w}^+ = \dot{w}^-, \quad \text{on } \Gamma. \quad (4.4)$$

Using methods from [17,18] we reduce boundary-value problem (4.4) to the following problem for analytic functions: find analytic function $\Phi^+(z)$ in G^+ and analytic function

$\Phi^-(z)$ in G^- satisfying the boundary-value condition $\Phi^+(s)=U(s)\Phi^-(s)$ on Γ , where $U(s) \neq 0$ on Γ , index $\kappa = \text{ind } U(s) = 0$, Φ^- has finite order on infinity $\Phi^-(\infty) = 0$ and $\Phi^+(z_0) = 0$ at the point $z_0 = x_0^1 + ix_0^2 \in G^+$. Therefore from [17, 18, 31] we obtain that the boundary-value problem (4.4) has only zero solution of class $C^{m-2,\nu}$. ($\Phi^+ \equiv 0, \Phi^- \equiv 0$).

For every admissible analytic in G^+ function Φ^+ the equation

$$\partial_{\bar{z}}\dot{w} + A\dot{w} + B\bar{\dot{w}} + E(\dot{w}) = \dot{\Psi}, \quad \text{in } G^+$$

has unique solution and is solved by method of successive approximations as (see [17, 18]):

$$\dot{w}_0(z) = \Phi^+(z) + T(-E(\dot{w}) + \dot{\Psi}),$$

$$\dot{w}_{k+1}(z) = \frac{1}{\pi} \int \int_{G^+} \frac{(A\dot{w}_k + B\bar{\dot{w}}_k)}{\zeta - z} d\xi d\eta + \Phi^+(z) + T(-E(\dot{w}) + \dot{\Psi}), \quad k = 0, 1, \dots,$$

where $\zeta = \xi + i\eta$,

$$T(f) = -\frac{1}{\pi} \int \int_{G^+} \frac{f(\zeta)}{\zeta - z} d\xi d\eta.$$

Then we have the following inequality: $\|\dot{w}\|_{m-2,\nu} \leq K_{20}(t)\|\dot{w}\|_{m-2,\nu}$ for all sufficiently small $t \geq 0$, where $K_{20}(t) < 1$.

Using methods from [17, 18, 31, 20] for finding the solutions of problem (4.3), theory of completely continuous operators and theory of Volterra operator equation, the theorem 1 from [30] we obtain that the problem (A') has only zero solution of class $C^{m-2,\nu}$ continuous by t .

The case 2) of theorem 1 is proved.

Proof of case 1) differs from proof of case 2) by the fact that $\dot{c}^+$ has one additional real parameter and Φ^+ is two-parametric function (with two real parameters).

References.

1. A.I. Bodrenko. On continuous almost ARG-deformations of hypersurfaces in Euclidean space [in Russian]. Dep. in VINITI 27.10.1992., N3084-T92, UDK 513.81, 14 pp.
2. A.I. Bodrenko. Some properties continuous ARG-deformations [in Russian]. Theses of international science conference "Lobachevskii and modern geometry Kazan, Kazan university publishing house, 1992 ., pp.15-16.
3. A.I. Bodrenko. On continuous ARG-deformations [in Russian]. Theses of reports on republican science and methodical conference, dedicated to the 200-th anniversary of N.I.Lobachevskii, Odessa, Odessa university publishing house, 1992 ., Part 1, pp.56-57.
4. A.I. Bodrenko. On extension of infinitesimal almost ARG-deformations closed hypersurfaces into analytic deformations in Euclidean spaces [in Russian]. Dep. in VINITI 15.03.1993., N2419-T93 UDK 513.81, 30 pp.

5. A.I. Bodrenko. On extension of infinitesimal almost ARG-deformations of hypersurface with boundary into analytic deformations [in Russian]. Collection works of young scholars of VolSU, Volgograd, Volgograd State University publishing house, 1993, pp.79-80.
6. A.I. Bodrenko. Some properties of continuous almost AR-deformations of hypersurfaces with prescribed change of Grassmannian image [in Russian]. Collection of science works of young scholars, Taganrog, Taganrog State Pedagogical Institute publishing house , 1994, pp. 113-120.
7. A.I. Bodrenko. On continuous almost AR-deformations with prescribed change of Grassmannian image [in Russian]. All-Russian school-colloquium on stochastic methods of geometry and analysis. Abrau-Durso. Publisher Moscow: "TVP". Theses of reports, 1994, pp. 15-16.
8. A.I. Bodrenko. Extension of infinitesimal almost ARG-deformations of hypersurfaces into analytic deformations [in Russian]. All-Russian school-colloquium on stochastic methods. Yoshkar-Ola. Publisher Moscow: "TVP". Theses of reports, 1995, pp. 24-25.
9. A.I. Bodrenko. Areal-recurrent deformations of hypersurfaces preserving Grassmannian image [in Russian]. Dissertation of candidate of physical-mathematical sciences. Novosibirsk, 1995, pp. 85.
10. A.I. Bodrenko. Areal-recurrent deformations of hypersurfaces preserving Grassmannian image [in Russian]. Author's summary of dissertation of candidate of physical-mathematical sciences. Novosibirsk, 1995, pp. 1-14.
11. A.I. Bodrenko. Some properties of ARG-deformations [in Russian]. Izvestiay Vuzov. Ser. Mathematics, 1996, N2, pp.16-19.
12. A.I. Bodrenko. Continuous almost ARG-deformations of surfaces with boundary [in Russian]. Modern geometry and theory of physical fields. International geometry seminar of N.I.Lobachevskii Theses of reports, Kazan, Publisher Kazan university, 1997, pp.20-21.
13. A.I. Bodrenko. Continuous almost AR-deformations of surfaces with prescribed change of Grassmannian image [in Russian]. Red. "Sib. mat. zhurnal. Sib. otd. RAN , Novosibirsk, Dep. in VINITI 13.04.1998., N1075-T98 UDK 513.81, 13 pp.
14. A.I. Bodrenko. Almost ARG-deformations of the second order of surfaces in Riemannian space [in Russian]. Surveys in Applied and Industrial Mathematics. 1998, Vol. 5, Issue 2, p.202. Publisher Moscow: "TVP".
15. A.I. Bodrenko. Almost *AR*-deformations of a surfaces with prescribed change of Grassmannian image with exterior connections [in Russian]. Red. zhurn. "Izvestiya vuzov. Mathematics. Kazan, Dep. in VINITI 03.08.1998, N2471 - B 98. P. 1-9.

16. A.I. Bodrenko. Properties of generalized G-deformations with areal condition of normal type in Riemannian space [in Russian]. Surveys in Applied and Industrial Mathematics. Vol. 7. Issue 2. (VII All-Russian school-colloquium on stochastic methods. Theses of reports.) P. 478. Moscow: TVP, 2000.
17. I.N. Vekua. Generalized Analytic Functions. Pergamon. New York. 1962.
18. I.N. Vekua. Generalized Analytic Functions [in Russian]. Moscow. Nauka. 1988.
19. I.N. Vekua. Some questions of the theory of differential equations and applications in mechanics [in Russian]. Moscow:"Nauka". 1991 . pp. 256.
20. A.V. Zabeglov. On decidability of one nonlinear boundary-value problem for AG-deformations of surfaces with boundary [in Russian]. Collection of science works. Transformations of surfaces, Riemannian spaces determined by given recurrent relations. Part 1. Taganrog. Taganrog State Pedagogical Institute publishing house. 1999. pp. 27-37.
21. V.T. Fomenko. On solution of the generalized Minkowski problem for surface with boundary [in Russian]. Collection of science works. Transformations of surfaces, Riemannian spaces determined by given recurrent relations. Part 1. Taganrog. Taganrog State Pedagogical Institute publishing house. 1999. pp. 56-65.
22. V.T. Fomenko. On uniqueness of solution of the generalized Christoffel problem for surfaces with boundary [in Russian]. Collection of science works. Transformations of surfaces, Riemannian spaces determined by given recurrent relations. Part 1. Taganrog. Taganrog State Pedagogical Institute publishing house. 1999. pp. 66-72.
23. V.T. Fomenko. On rigidity of surfaces with boundary in Riemannian space [in Russian]. Doklady Akad. Nauk SSSR. 1969 . Vol. 187, N 2, pp. 280-283.
24. V.T. Fomenko. ARG-deformations of hypersurfaces in Riemannian space [in Russian]. //Dep. in VINITI 16.11.1990 N5805-B90
25. S.B. Klimentov. On one method of construction the solutions of boundary-value problems in the bending theory of surfaces of positive curvature [in Russian]. Ukrainian geometry sbornik. pp. 56-82.
26. M.A. Krasnoselskii. Topological methods in the theory of nonlinear problems [in Russian]. Moscow, 1965.
27. L.P. Eisenhart. Riemannian geometry [in Russian]. Izd. in. lit., Moscow 1948. (Eisenhart Luther Pfahler. Riemannian geometry. 1926.)
28. J.A. Schouten, D.J. Struik. Introduction into new methods of differential geometry [in Russian]. Volume 2. Moscow. GIIL. 1948 . (von J.A. Schouten und D.J. Struik. Einführung in die neueren methoden der differentialgeometrie. Zweite vollständig umgearbeitete Auflage. Zweiter band. 1938.)

29. I. Kh. Sabitov. //VINITI. Results of science and technics. Modern problems of mathematics [in Russian]. Fundamental directions. Vol.48, pp.196-271.
30. Andrei I. Bodrenko. The solution of the Minkowski problem for open surfaces in Riemannian space. Preprint 2007. arXiv:0708.3929v1 [math.DG]
31. F.D. Gakhov. Boundary-value problems [in Russian]. GIFML. Moscow. 1963.